Product Datasheet

5-Lipoxygenase Antibody - BSA Free NB110-58749

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB110-58749

Updated 12/20/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB110-58749




NB110-58749

5-Lipoxygenase Antibody - BSA Free

5-Lipoxygenase Antibody - BSA Free	
Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 30% Glycerol
Target Molecular Weight	80 kDa
Product Description	
Host	Rabbit
Gene ID	240
Gene Symbol	ALOX5
Species	Human, Mouse, Rat, Bovine, Rabbit
Immunogen	A synthetic peptide made to a C-terminal region within residues 600-674 of human 5-Lipoxygenase. [Swiss-Prot# P09917]
Product Application Details	
Applications	Western Blot, Simple Western
Recommended Dilutions	Western Blot 1:500-1:1000, Simple Western 10 ug/ml
Application Notes	This 5 Lipoxygenase antibody is useful for Western Blot, where a band is seen at approx. 80 kDa. There may be some non-specific bands at non-interfering lower molecular weights.
	In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. Separated by Size-Wes, Sally Sue/Peggy Sue. The observed molecular weight of the protein may vary from the listed predicted melocular weight due to post translational medifications, post translations.

molecular weight due to post translational modifications, post translation

cleavages, relative charges, and other experimental factors.



Publications

Di Federico M, Ancora M, Luciani M et al. Pro-Inflammatory Response of Bovine Polymorphonuclear Cells Induced by Mycoplasma mycoides subsp. mycoides Front Vet Sci 2020-03-27 [PMID: 32292794] (WB, Bovine)



Procedures

Protocol specific for 5 Lipoxygenase Antibody (NB110-58749)

5-Lipoxygenase Antibody:

Western Blot Protocol

- 1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 38 ug of total protein per lane.
- 2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
- 3. Rinse membrane with dH2O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
- 4. Rinse the blot in TBS for approximately 5 minutes.
- 5. Block the membrane using 5% non-fat dry milk + 1% BSA in TBS, 1 hour at room temperature.
- 6. Rinse the membrane in dH2O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
- 7. Dilute the rabbit anti-LOX5 primary antibody (NB 110-58749) in blocking buffer and incubate 2 hours at room temperature.
- 8. Rinse the membrane in dH2O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
- 9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
- 10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
- 11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB110-58749

NB820-59174 Blood Leukocyte Whole Cell Lysate

NB110-58749PEP 5-Lipoxygenase Antibody Blocking Peptide

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-58749

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

