# **Product Datasheet**

## Testin Antibody NBP1-87987

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.



#### **Publications: 3**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-87987

Updated 8/22/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-87987




## NBP1-87987

Testin Antibody

Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol
Product Description	
Host	Rabbit
Gene ID	26136
Gene Symbol	TES
Species	Human
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non- tested species: Mouse (82%), Rat (85%). Reactivity reported in scientific literature (PMID: 23227862)
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: QLPAHDQDPSKCHELSPREVKEMEQFVKKYKSEALGVGDVKLPCEMDAQGPK QMNIPGGDRSTPAAVGAMEDKSAEHKRTQYSCYCCKLSMKEGDPAIYAERAG YDKLWHPACFVCSTCHELLVD
Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Simple Western reported by internal validation, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:200
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. Separated by Size-Wes, Sally Sue/Peggy Sue. Use in IHC-P reported in scientific literature (PMID:34440214).







	Page 3 of 6 V.20.1 Updated 8/22/2023
Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human testis shows moderate cytoplasmic positivity in cells in seminiferous ducts.	
Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human endometrium moderate cytoplasmic positivity in glandular cells.	
Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human liver shows no cytoplasmic positivity in hepatocytes as expected.	
Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human endometrium shows moderate cytoplasmic positivity in glandular cells.	



Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human urinary bladder shows strong cytoplasmic positivity in urothelial cells. Immunohistochemistry-Paraffin: Testin Antibody [NBP1-87987] - Staining of human liver shows no positivity in hepatocytes as expected. Simple Western: Testin Antibody [NBP1-87987] - Simple Western lane view shows a specific band for TES in 0.2 mg/ml of RT-4 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system. Simple Western: Testin Antibody [NBP1-87987] - Electropherogram image(s) of corresponding Simple Western lane view. Testin antibody was used at 1:20 dilution on RT-4 lysate(s). 40 MW (kDa)





#### **Publications**

Popiel-Kopaczyk A, Grzegrzolka J, Piotrowska A et al. The Expression of Testin, Ki-67 and p16 in Cervical Cancer Diagnostics Current issues in molecular biology 2023-01-05 [PMID: 36661518] (IHC-P, Human)

Details:

Dilution used in IHC-P 1:400

Popiel A, Piotrowska A, Sputa-Grzegrzolka P Et al. Preliminary Study on the Expression of Testin, p16 and Ki-67 in the Cervical Intraepithelial Neoplasia Biomedicines 2021-08-13 [PMID: 34440214] (IHC-P, Human)

Danielsson F, Wiking M, Mahdessian D et al. RNA deep sequencing as a tool for selection of cell lines for systematic subcellular localization of all human proteins. J Proteome Res 2013-01-04 [PMID: 23227862]





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP1-87987

NBP1-87987PEP	Testin Recombinant Protein Antigen
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-87987

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

