# **Product Datasheet**

# TOP2B Antibody NBP1-89527

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-89527

Updated 3/12/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-89527



# NBP1-89527


**TOP2B** Antibody

TOP2B Antibody	
Product Information	
0.1 ml	
Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Polyclonal	
0.02% Sodium Azide	
IgG	
Immunogen affinity purified	
PBS (pH 7.2) and 40% Glycerol	
Product Description	
Rabbit	
7155	
TOP2B	
Human, Mouse, Rat	
Human reactivity reported in scientific literature (PMID: 26439921).	
This antibody was developed against Recombinant Protein corresponding to amino acids: LWKEDLAAFVEELDKVESQEREDVLAGMSGKAIKGKVGKPKVKKLQLEETMPS PYGRRIIPEITAMKADASKKLLKKKKGDLDTAAVKVEFDEEFSGAPVEGAGEEA LTPSVPINKGPKPKREKKEPGTRVRKTPTSSGKPSA	
Product Application Details	
Western Blot, Simple Western, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockdown Validated	
Western Blot 0.04-0.4 ug/ml, Simple Western 1:100, Immunohistochemistry 1:500 - 1:1000, Immunocytochemistry/Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:500 - 1:1000, Knockdown Validated	
For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.  In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. Separated by Size-Wes, Sally Sue/Peggy Sue.	



## **Images**


Western Blot: TOP2B Antibody [NBP1-89527] - Analysis in U2OS cells transfected with control siRNA, target specific siRNA probe #1 and #2. Remaining relative intensity is presented. Loading control: Anti-GAPDH.




Immunocytochemistry/Immunofluorescence: TOP2B Antibody [NBP1-89527] - Staining of human cell line U-251 MG shows localization to nucleoplasm. Antibody staining is shown in green.



Immunohistochemistry-Paraffin: TOP2B Antibody [NBP1-89527] - Staining of human pancreas shows weak to moderate nuclear positivity in islets of Langerhans and exocrine glandular cells.




Western Blot: TOP2B Antibody [NBP1-89527] - Analysis in mouse cell line NIH-3T3 and rat cell line NBT-II.



Page 3 of 5 v.20.1 Updated 3/12/2023 Immunohistochemistry-Paraffin: TOP2B Antibody [NBP1-89527] -Staining of human cerebral cortex shows strong nuclear positivity in neurons. Immunohistochemistry-Paraffin: TOP2B Antibody [NBP1-89527] -Staining of human gastrointestinal shows strong nuclear positivity in glandular cells. Immunohistochemistry-Paraffin: TOP2B Antibody [NBP1-89527] -Staining of human kidney shows moderate nuclear positivity in cells in glomeruli and tubules. Simple Western: TOP2B Antibody [NBP1-89527] - Simple Western lane view shows a specific band for TOP2B in 0.2 mg/ml of RT-4 lysate. This experiment was performed under reducing conditions using the 66-440 kDa separation system.



Simple Western: TOP2B Antibody [NBP1-89527] - Electropherogram image(s) of corresponding Simple Western lane view. TOP2B antibody was used at 1:100 dilution on RT-4 lysate(s).



#### **Publications**

Wang T, Kyle AH, Baker JHE et al. DNA-PK inhibition extends the therapeutic effects of Top2 poisoning to non-proliferating cells, increasing activity at a cost Scientific Reports 2023-08-01 [PMID: 37528151] (IHC-Fr, IHC)

Lee S, Jun YW, Linares GR et al. Downregulation of Hsp90 and the antimicrobial peptide Mtk suppresses poly(GR)-induced neurotoxicity in C9ORF72-ALS/FTD Neuron 2023-03-10 [PMID: 36931278] (WB, Drosophila)

#### Details:

Dilution used in WB 1:2,000

Hu HL, Shiflett LA, Kobayashi M et al. TOP2b-Dependent Nuclear DNA Damage Shapes Extracellular Growth Factor Responses via Dynamic AKT Phosphorylation to Control Virus Latency Mol. Cell 2019-03-26 [PMID: 30930055] (WB, Rat)

Ali A, Biswas A, Pal M. HSF1 mediated TNF-a production during proteotoxic stress response pioneers proinflammatory signal in human cells. FASEB J. 2018-10-11 [PMID: 30307772] (WB, Human)

Aviner R, Shenoy A, Elroy-Stein O, Geiger T. Uncovering Hidden Layers of Cell Cycle Regulation through Integrative Multi-omic Analysis. PLoS Genet 2015-10-01 [PMID: 26439921] (WB, Human)





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP1-89527**

NBP1-89527PEP TOP2B Recombinant Protein Antigen

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-89527

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

