

Product Datasheet

PKM1 Antibody - BSA Free

NBP2-14833

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 3 **Publications: 18**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-14833

Updated 12/20/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-14833



NBP2-14833


PKM1 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.17 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 30% Glycerol
Product Description	
Host	Rabbit
Gene ID	5315
Gene Symbol	PKM
Species	Human, Mouse, Bovine
Reactivity Notes	Bovine reactivity reported in scientific literature (PMID: 25416385).
Specificity/Sensitivity	This antibody is specific for the M1 isoform of PKM.
Immunogen	A synthetic peptide made to a C-terminal portion of the human PKM1 protein (between residues 350-450) [UniProt P14618-2].
Product Application Details	
Applications	Western Blot, Simple Western, Immunoblotting, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 0.5 ug/ml, Simple Western 1:400, Immunohistochemistry 1:400, Immunocytochemistry/ Immunofluorescence reported in multiple pieces of scientific literature, Immunohistochemistry-Paraffin 1:400, Immunoblotting reported in scientific literature (PMID 25416385)
Application Notes	<p>In Western blot a band is seen ~57 kDa in human and mouse skeletal muscle, brain and heart. In IHC-P, staining was observed in the cytoplasm of human breast cancer tissue. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




Images


Western Blot: PKM1 Antibody [NBP2-14833] - Expression profile of PKM isoforms in tissues from mouse organs. PKM1 and PKM2 were detected by Western blotting in under the same experimental conditions as Figure 1a at the same time. The full-length blots are presented in Supplementary Figure S2b. Results are presented as the mean \pm SD (** P < 0.01). Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/srep08647>), licensed under a CC-BY license.




Immunohistochemistry-Paraffin: PKM1 Antibody [NBP2-14833] - PKM1 antibody was tested in human breast cancer using DAB with hematoxylin counterstain.




Western Blot: PKM1 Antibody [NBP2-14833] - PKM1 antibody was tested in 1. human skeletal muscle 2. mouse skeletal muscle 3. human brain 4. mouse brain and 5. human heart cell lysate.




Western Blot: PKM1 Antibody [NBP2-14833] - Detection of PKM1 on HEK293T and U-87 cells. Lane 1: HEK293T cells (major isoform is PKM2, human). Lane 2: HEK293T cells with plasmid overexpressing mouse PKM1. Lane 3: HEK293T cells with plasmid overexpressing mouse PKM2. Lane 4: U-87 cells with control shRNA (major isoform is PKM2, human). Lane 5: U-87 cells with PTB/A1/A2 shRNAs (major isoform is PKM1, human). Photo courtesy of product review by verified customer.




Western Blot: PKM1 Antibody [NBP2-14833] - Expression of PTB1, PKM1, and PKM2 proteins at 72 h after transfection of DLD-1, RD or KYM-1 cells with miR-133b (10, 20 nM). Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/srep08647>), licensed under a CC-BY license.



Western Blot: PKM1 Antibody [NBP2-14833] - Expression profile of PKM isoforms in various cancer cell lines, control cell lines, and cells in primary culture. PKM1 and PKM2 were detected by Western blotting under the same experimental conditions at the same time. The full-length blots are presented in Supplementary Figure S2a. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/srep08647>), licensed under a CC-BY license.



Simple Western: PKM1 Antibody [NBP2-14833] - Simple Western lane view shows a specific band for PKM1 in 0.5 mg/ml of Human Brain lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

Li Y, Chen X, Huang H et al. A feedback loop between NONHSAT024276 and PTBP1 inhibits tumor progression and glycolysis in HCC by increasing the PKM1/PKM2 ratio *Cancer Science* 2023-04-01 [PMID: 36529521] (WB)

Deck M, Van Hameren G, Campbell G et al. Physiology of PNS axons relies on glycolytic metabolism in myelinating Schwann cells *PLoS one* 2022-10-04 [PMID: 36194565] (WB, IHC-Fr, Mouse)

Han J, Hyun J, Park J et al. Aberrant role of pyruvate kinase M2 in the regulation of gamma-secretase and memory deficits in Alzheimer's disease *Cell reports* 2021-12-07 [PMID: 34879266]

Tian, L, Xiong, P Y Et al. Supra-coronary aortic banding improves right ventricular function in experimental pulmonary arterial hypertension in rats by increasing systolic right coronary artery perfusion. *Acta Physiol (Oxf)* 2020-08-01 [PMID: 32339403] (ICC/IF, Human)

Zhang H, Wang D et al. Metabolic and Proliferative State of Vascular Adventitial Fibroblasts in Pulmonary Hypertension Is Regulated Through a MicroRNA-124/PTBP1 (Polypyrimidine Tract Binding Protein 1)/Pyruvate Kinase Muscle Axis. *Circulation* 2017-12-19 [PMID: 28972001] (WB, Bovine)

Okazaki Mitsuyoshi, Fushida Sachio, Tsukada Tomoya et al. The effect of HIF-1a and PKM1 expression on acquisition of chemoresistance. *Cancer Management and Research* 2018-01-01 [PMID: 30013393] (WB, IF/IHC, Mouse)

Shanmugasundaram K, Nayak BK, Friedrichs WE et al. NOX4 functions as a mitochondrial energetic sensor coupling cancer metabolic reprogramming to drug resistance. *Nat Commun.* 2017-10-19 [PMID: 29051480] (WB, Human)

Minami K, Taniguchi K, Sugito N et al. MiR-145 negatively regulates Warburg effect by silencing KLF4 and PTBP1 in bladder cancer cells. *Oncotarget.* 2017-05-16 [PMID: 28380435] (WB, Human)

Takai T, Yoshikawa Y, Inamoto T et al. A Novel Combination RNAi toward Warburg Effect by Replacement with miR-145 and Silencing of PTBP1 Induces Apoptotic Cell Death in Bladder Cancer Cells *Int J Mol Sci* 2017-01-17 [PMID: 28106737] (WB, Human)

Sugiyama T, Taniguchi K, Matsushashi N et al. MiR-133b inhibits growth of human gastric cancer cells by silencing pyruvate kinase muscle-splicer polypyrimidine tract-binding protein 1. *Cancer Sci.* 2016-12-01 [PMID: 27696637] (WB, Human)

J Casson R, P M Wood J, Han G et al. M-Type Pyruvate Kinase Isoforms and Lactate Dehydrogenase A in the Mammalian Retina: Metabolic Implications. *Invest. Ophthalmol. Vis. Sci.* 2016-01-01 [PMID: 26780311] (WB, IHC-P, Mouse)

Taniguchi K, Sugito N, Kumazaki M et al. Positive feedback of DDX6/c-Myc/PTBP1 regulated by miR-124 contributes to maintenance of the Warburg effect in colon cancer cells *Biochim. Biophys. Acta.* 2015-07-02 [PMID: 26144048] (WB, Human)

Details:

PKM2 antibody was used for WB assay on lysates of human colon cancer cells (DLD-1 cells and WiDr cells) transfected with miR-124 or siR-PTBP1 (Fig. 3B and 3C). WB was also performed on lysates after the transfection of colon cancer cells with siR-c-Myc or siR-DDX6 (Fig. 4A and 4B), and on DLD-1 cells/WiDr cells that were subjected to PKM1 and/or PKM2 knockdown using siRNAPKM1 and siRNAPKM2 (Fig. 5B and 5E).

More publications at <http://www.novusbio.com/NBP2-14833>

Procedures

Western Blot protocol for PKM1 Antibody (NBP2-14833)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 25 ug of total protein per lane.
2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot.
5. Block the membrane using standard blocking buffer for at least 1 hour.
6. Wash the membrane in wash buffer three times for 10 minutes each.
7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
8. Wash the membrane in wash buffer three times for 10 minutes each.
9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

*Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

Immunohistochemistry-Paraffin protocol for PKM1 Antibody (NBP2-14833)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-14833

NB820-59253	Human Skeletal Muscle Whole Tissue Lysate (Adult Whole Normal)
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-14833

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

