

Product Datasheet

BAG3 Antibody NBP2-27398

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 13

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-27398

Updated 4/21/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-27398



NBP2-27398

BAG3 Antibody

Product Information	
Unit Size	0.05 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Unpurified
Buffer	50 ul neat whole antisera

Product Description	
Host	Rabbit
Gene Symbol	BAG3
Species	Human, Mouse, Rat
Reactivity Notes	Use in Rat reported in scientific literature (PMID:32132902).
Immunogen	A recombinant protein fragment corresponding to the C-terminal 196 amino acids of human BAG-3. Bag-3


Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:2000~, Simple Western 1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:200, Immunoprecipitation 1:50-1:200, Immunohistochemistry-Paraffin 1:1000-1:5000~, Immunohistochemistry-Frozen 1:1000-1:5000
Application Notes	In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. Separated by Size-Wes, Sally Sue/Peggy Sue.

Images


Simple Western: BAG3 Antibody [NBP2-27398] - Simple Western lane view shows a specific band for BAG3 in 0.5 mg/ml of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Immunohistochemistry: BAG3 Antibody [NBP2-27398] - IHC of BAG3 in IR myocardium. BAG3 expression is significantly decreased in the STZ group than those in the CON group. BAG3 expression is also significantly reduced in the IR CON and IR STZ groups as compared with the CON group. IR STZ hearts further depress BAG3 expression than IR CON hearts. Data are expressed as mean \pm SEM (n = 3) using the single values. * p < 0.05 vs. CON, a p < 0.05 vs. STZ, b p < 0.05 vs. IR CON. Image collected and cropped by CiteAb from the following publication (<https://www.mdpi.com/2076-3921/9/8/679>) licensed under a CC-BY license.




Western Blot: BAG3 Antibody [NBP2-27398] - Western blot of BAG3 in IR myocardium. BAG3 expression is significantly decreased in the STZ group than those in the CON group. BAG3 expression is also significantly reduced in the IR CON and IR STZ groups as compared with the CON group. IR STZ hearts further depress BAG3 expression than IR CON hearts. Data are expressed as mean \pm SEM (n = 3) using the single values. * p < 0.05 vs. CON, a p < 0.05 vs. STZ, b p < 0.05 vs. IR CON. Image collected and cropped by CiteAb from the following publication (<https://www.mdpi.com/2076-3921/9/8/679>) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: BAG3 Antibody [NBP2-27398] - Antibody was tested in A431 cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red).




Western Blot: BAG3 Antibody [NBP2-27398] - Analysis of BAG3 using NBP2-27398 at 1:2000. Tissue lysates, normalized for total protein (20 ug/lane), were from a 4 week old male mouse. BAG-3 expression was detected at highest levels in skeletal (quadriceps and diaphragm) and smooth (heart) muscle specimens.



Immunohistochemistry-Frozen: BAG3 Antibody [NBP2-27398] - Frozen mouse muscle tissue section stained for BAG-3 expression using NBP2-27398 at 1:2000. The tissue section was fixed in 3.8% paraformaldehyde prior to staining. BAG-3 localizes with Z-disk proteins.



Mesenteric vascular Western blot graph (A) and analysis of 4-HNE (B), CaSR (C), p-eNOS (D) and BAG3 (E) are demonstrated. Microvascular reactivity from mesenteric arterioles to a vasoconstrictor (norepinephrine, NE, F) and vasodilator (acetylcholine, Ach, G) is displayed in CON and STZ groups. All data are presented as the mean \pm SEM (n = 3) using the single values in each test. * p < 0.05 vs. CON group.



Publications

Lu S, Hu J, Arogundade OA et al. Heat-shock chaperone HSPB1 regulates cytoplasmic TDP-43 phase separation and liquid-to-gel transition *Nature Cell Biology* 2022-09-08 [PMID: 36075972]

Damiani V, Lamolinara A, Cicalini I et al. Pancreatic beta-cell specific BAG3 knockout results in chronic hyperinsulinemia inducing insulin resistance *Molecular metabolism* 2023-06-10 [PMID: 37308077] (ICC/IF, Mouse)

Damiani V, Cufaro MC, Fucito M et al. Proteomics Approach Highlights Early Changes in Human Fibroblasts-Pancreatic Ductal Adenocarcinoma Cells Crosstalk Cells 2022-03-29 [PMID: 35406724] (WB, Human)

Mukherjee, T, Ramaglia, V Et al. The eIF2 alpha kinase HRI triggers the autophagic clearance of cytosolic protein aggregates. *J Biol Chem* 2020-12-02 [PMID: 33410405] (WB, Mouse)

Li X, Lin G, Liu T Et al. Postnatal development of BAG3 expression in mouse cerebral cortex and hippocampus *Brain structure & function* 2021-08-06 [PMID: 34357438]

Yapa Abeywardana M, Samarasinghe KTG, Munkanatta Godage D, Ahn YH Identification and Quantification of Glutathionylated Cysteines under Ischemic Stress *Journal of proteome research* 2021-08-12 [PMID: 34382403]

Chien CY, Wen TJ, Cheng YH et al. Diabetes Upregulates Oxidative Stress and Downregulates Cardiac Protection to Exacerbate Myocardial Ischemia/Reperfusion Injury in Rats *Antioxidants (Basel)* 2020-07-29 [PMID: 32751309] (IHC-P, WB, Rat)

Ormeno F, Hormazabal J, Moreno J et Al. Chaperone Mediated Autophagy Degrades TDP-43 Protein and Is Affected by TDP-43 Aggregation *Front Mol Neurosci* 2020-02-18 [PMID: 32132902] (WB, Human, Rat)

Li X, Lu J, Kan Q et al. Metabolic reprogramming is associated with flavopiridol resistance in prostate cancer DU145 cells *Sci Rep* 2017-07-11 [PMID: 28698547] (WB, Human)

Liao Q, Ozawa F, Friess H et al. The anti-apoptotic protein BAG-3 is overexpressed in pancreatic cancer and induced by heat stress in pancreatic cancer cell lines. *FEBS Lett.* 2001-08-17 [PMID: 11513873] (IHC-P, WB, Human)

Details:

WB: human tissues and cell lines (Figs 1, 4-6); IHC (P): human pancreas tissue (Fig 2).

Aust Stefanie, Pils Sophie, Polterauer Stefan et al. Expression of Bcl-2 and the antiapoptotic BAG family proteins in ovarian cancer. *Appl Immunohistochem Mol Morphol.* 2013-12-01 [PMID: 23455184] (IHC-P, Human)

Goldman A, Chen HD, Roesly HB et al. Characterization of squamous esophageal cells resistant to bile acids at acidic pH: implication for Barrett's esophagus pathogenesis. *Am J Physiol Gastrointest Liver Physiol.* 2011-02-01 [PMID: 21127259] (WB, Human)

Details:

BAG-3 (IMG-5676). WB: Normal human esophageal epithelial cell lines (HET1A & HET1AR), Fig 4D.

More publications at <http://www.novusbio.com/NBP2-27398>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-27398

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-72276-50ug	Recombinant Human BAG3 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-27398

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

