Product Datasheet

Cytokeratin 18 Antibody (DC10) NBP2-29461

Unit Size: 0.1 mg

Store at 4C.



Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-29461

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-29461



NBP2-29461

Cytokeratin 18 Antibody (DC10)

0.1 mg
0.2 mg/ml
Store at 4C.
Monoclonal
DC10
0.05% Sodium Azide
IgG1 Kappa
Protein A or G purified
10 mM PBS with 0.05% BSA
45 kDa
200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-33082) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Mouse
3875
KRT18
Human, Ferret (Negative), Bovine (Negative), Canine (Negative), Hamster (Negative), Mouse (Negative), Porcine (Negative), Rat (Negative), Sheep (Negative)
Does not react with Mouse, Rat, Ferret, Hamster, Sheep, Bovine, Canine or Porcine.
Epithelial Marker
This monoclonal antibody reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs to the type A (acidic) subfamily of low molecular weight keratins, exists in combination with cytokeratin 8. It is reported that tissues from gastrointestinal tract are positive for both cytokeratin 8 and 18 but do not contain cytokeratin 14. Tissues from gastrointestinal tract, respiratory tract and urogenital tract, as well as endocrine and exocrine tissues and mesothelial cells are positive for cytokeratin 18.
PMC-42 human breast carcinoma cells
Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Western Blot 1-2 ug/ml, Simple Western 10 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry 0.5-1ug/ml, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml




Application Notes


Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined. In Simple Western only 10 - 15 ul of the recommended dilution is used per data point. Separated by Size.

Images

cells.

Western Blot: Cytokeratin 18 Antibody (DC10) [NBP2-29461] - Western Blot Analysis of human HCT116 cell lysate using Cytokeratin 18 Antibody (DC10).





Western Blot: Cytokeratin 18 Antibody (DC10) [NBP2-29461] - Analysis using the Azide and BSA Free version of NBP2-29461. Detection of Keratin 18 in 1) HeLa and 2) A431 lysate using Keratin 18 antibody at 1 ug/ml. goat anti-mouse Ig HRP secondary antibody and PicoTect ECL


substrate solution were used for this test.

(kDa) 200 —	1	2
¹¹⁶ 97 ==		
66 — 55 —		
36 —	٦	•
31 —		
21 —		
14 —		

MANA/







Publications

Soeta K, Iuchi K, Hisatomi H, Yokoyama C Generation of Rat Monoclonal Antibody for Cytokeratin 18 by Immunization of Three-Dimensional-Cultured Cancer Cells Monoclon Antib Immunodiagn Immunother 2020-10-16 [PMID: 33064594]

Yoon HJ, Jo BC, Shin WJ et al. Comparative immunohistochemical study of ameloblastoma and ameloblastic carcinoma. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2011-12-01 [PMID: 22014999]

Terada T. Angiosarcoma of the oral cavity. Head Neck Pathol. 2011-03-01 [PMID: 20924728]

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-29461

H00003875-P01-10ug	Recombinant Human Cytokeratin 18 GST (N-Term) Protein
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-29461

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

